

AMS 3 formula sheet

All formula with an asterisk must be known by heart. All others are optional and will be given to you in exams if required.

1 Lines

The equation of a line with slope m and y-intercept b is

$$y = f(x) = mx + b \quad (*) \quad (1)$$

If the line goes through 2 points A and B with coordinates (x_A, y_A) and (x_B, y_B) then

$$m = \frac{y_B - y_A}{x_B - x_A} \quad (*) \quad (2)$$

2 Quadratic equations

$$y = f(x) = ax^2 + bx + c \quad (*) \quad (3)$$

The graph of $y = f(x)$ is a parabola. It has a minimum (i.e. parabola opens upwards) if $a > 0$. It has a maximum (i.e. parabola opens downwards) if $a < 0$. The minimum/maximum is at the location x_V with

$$x_V = -\frac{b}{2a} \quad (*) \quad (4)$$

It has roots (i.e it intersects the x -axis) when $y = f(x) = 0$. The solutions to this equation depends on the value of D :

$$D = b^2 - 4ac \quad (*) \quad (5)$$

- if $D < 0$ there are no solutions. The parabola does not intercept the x -axis. The function $f(x) = ax^2 + bx + c$ cannot be factored.
- if $D = 0$ there is one solution. The parabola just touches the x -axis at the point

$$x_1 = x_V = -\frac{b}{2a} \quad (*) \quad (6)$$

The function $f(x) = ax^2 + bx + c$ is factored as

$$f(x) = a(x - x_1)^2 \quad (*) \quad (7)$$

- if $D > 0$ there are two solutions. The parabola intercepts the x -axis in the two points

$$x_1 = \frac{-b - \sqrt{D}}{2a}, x_2 = \frac{-b + \sqrt{D}}{2a} \quad (*) \quad (8)$$

The function $f(x) = ax^2 + bx + c$ is factored as

$$f(x) = a(x - x_1)(x - x_2) \quad (*) \quad (9)$$

3 Polynomial functions

$$y = f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n \quad (*) \quad (10)$$

a_n is the leading coefficient, n is the order of the polynomial.

The factored form of f is

$$f(x) = a_n(x - x_1)(x - x_2)(x - x_3) \dots (x - x_m)q(x) \quad (*) \quad (11)$$

where x_i are all possible solutions to $f(x) = 0$ and $q(x)$ is a polynomial of order $n - m$, leading coefficient 1, with no roots ($q(x) \neq 0$).

4 Rational functions

$$y = f(x) = \frac{p(x)}{q(x)} \quad (*) \quad (12)$$

where $p(x)$ and $q(x)$ are polynomial functions.

The roots of $p(x)$ are the roots of $f(x)$. The roots of $q(x)$ are the asymptotes of $f(x)$.

5 Power functions

$$y = f(x) = x^a \quad (*) \quad (13)$$

Properties:

$$x^{a+b} = x^a x^b \quad (*) \quad (14)$$

$$x^{-a} = \frac{1}{x^a} \quad (*) \quad (15)$$

$$x^{a-b} = \frac{x^a}{x^b} \quad (*) \quad (16)$$

$$x^{ab} = (x^a)^b = (x^b)^a \quad (*) \quad (17)$$

6 Exponential functions

Exponential in base a :

$$y = f(x) = a^x \text{ with } a > 0 \quad (*) \quad (18)$$

Natural exponential (exponential in base e with $e = 2.71828 \dots$):

$$y = f(x) = e^x = \exp(x) \quad (*) \quad (19)$$

Properties of all exponential functions:

$$a^{x+z} = a^x a^z \quad (*) \quad (20)$$

$$a^{-x} = \frac{1}{a^x} \quad (*) \quad (21)$$

$$a^{x-z} = \frac{a^x}{a^z} \quad (*) \quad (22)$$

$$a^{xz} = (a^x)^z = (a^z)^x \quad (*) \quad (23)$$

7 Logarithmic functions

Logarithm in base a is the inverse of the exponential in base a :

$$y = \log_a(x) \text{ is equivalent to } x = a^y \quad (*) \quad (24)$$

Natural logarithm (logarithm in base e) is the inverse of the natural logarithm:

$$y = \log_e(x) = \ln(x) \text{ is equivalent to } x = e^y \quad (*) \quad (25)$$

Inverse relations:

$$\log_a(a^x) = x \quad (*) \quad (26)$$

$$a^{\log_a(x)} = x \quad (*) \quad (27)$$

$$\ln(e^x) = x \quad (*) \quad (28)$$

$$e^{\ln(x)} = x \quad (*) \quad (29)$$

Properties of all logarithmic functions (where a is a positive constant).

$$\log_a(xy) = \log_a(x) + \log_a(y) \quad (*) \quad (30)$$

$$\log_a\left(\frac{1}{x}\right) = -\log_a(x) \quad (*) \quad (31)$$

$$\log_a\left(\frac{x}{y}\right) = \log_a(x) - \log_a(y) \quad (*) \quad (32)$$

$$\log_a(x^c) = c \log_a(x) \quad (*) \quad (33)$$

Relations for changing bases:

- From an exponential function in base a to the natural exponential:

$$a^x = e^{x \ln a} \quad (*) \quad (34)$$

- From a logarithmic function in base a to the natural logarithm:

$$\log_a(x) = \frac{\ln x}{\ln a} \quad (*) \quad (35)$$

8 Trigonometric functions

The basic trigonometric functions are:

$$y = f(x) = \sin(x) \quad (*) \quad (36)$$

$$y = f(x) = \cos(x) \quad (*) \quad (37)$$

$$y = f(x) = \tan(x) = \frac{\sin(x)}{\cos(x)} \quad (*) \quad (38)$$

Table of values you have you know (*):

Angle (degree)	Angle (radian)	sin	cos	tan
0	0	0	1	0
30	$\pi/6$	0.5	$\sqrt{3}/2$	$1/\sqrt{3}$
45	$\pi/4$	$\sqrt{2}/2$	$\sqrt{2}/2$	1
60	$\pi/3$	$\sqrt{3}/2$	0.5	$\sqrt{3}$
90	$\pi/2$	1	0	not defined

Properties:

$$\cos^2 x + \sin^2 x = 1 \quad (*) \quad (39)$$

$$\sin(2x) = 2 \sin x \cos x \quad (40)$$

$$\cos(2x) = \cos^2 x - \sin^2 x \quad (41)$$

$$\tan(2x) = \frac{2 \tan x}{1 - \tan^2 x} \quad (42)$$

Other addition/multiplication formula

$$\cos(a + b) = \cos a \cos b - \sin a \sin b \quad (43)$$

$$\cos(a - b) = \cos a \cos b + \sin a \sin b \quad (44)$$

$$\sin(a + b) = \sin a \cos b + \cos a \sin b \quad (45)$$

$$\sin(a - b) = \sin a \cos b - \cos a \sin b \quad (46)$$

$$\tan(a + b) = \frac{\tan a + \tan b}{1 - \tan a \tan b} \quad (47)$$

$$\tan(a - b) = \frac{\tan a - \tan b}{1 + \tan a \tan b} \quad (48)$$