

CHAPTER 3 Second order linear PDEs - Canonical form

3.1 Definition

- A second order linear PDE has the general form

$$\begin{aligned} L(u) = & \quad a(x,y)u_{xx} + 2b(x,y)u_{xy} + c(x,y)u_{yy} \quad \leftarrow \text{principal part} \\ & + d(x,y)u_x + e(x,y)u_y + f(x,y)u = g(x,y) \end{aligned}$$

- The principal part is the part of the equation that only involves second-order derivatives.
- The discriminant of the linear operator L is defined as

$$\delta(\lambda) = b^2(x,y) - a(x,y)c(x,y)$$

and, in the general case, will be a function of x and y .

Examples

- the wave equation: $u_{tt} = c^2 u_{xx}$

write as $u_{tt} - c^2 u_{xx} = 0$

so $\delta(\lambda) = c^2(x)$

- the heat equation $u_t = k u_{xx}$

write as $k u_{xx} - u_t = 0$

so $\delta(\lambda) = -k \cdot 0 = 0$

• the Laplace equation - $u_{xx} + u_{yy} = 0$

$$\rightarrow \delta(x) = -1$$

Definition: An operator is hyperbolic/parabolic/elliptic at a point (x, y) if $\delta(x)$ is respectively >0 , $=0$ or <0 at this point.

The operator for the wave equation is hyperbolic at all points (assume $c^2(x) > 0$).

_____ the heat equation is parabolic at all points

_____ Laplace equation is elliptic at all points

\Rightarrow An equation is hyperbolic/parabolic/elliptic in a domain D if its corresponding operator is hyperbolic/parabolic/elliptic at all points in D .

3.2 Properties of the discriminant under a change of coordinate

The sign of the discriminant of an operator L is invariant under a change of coordinates from (x, y) to (ξ, η) (such that the Jacobian $J = \xi_x \eta_y - \xi_y \eta_x \neq 0$ for all (x, y)).

In other words, the type of an equation is an intrinsic property of the equation and is independent of the coordinate system in which the equation is written.

Proof

Let

$$\xi = \xi(x, y)$$

$$\eta = \eta(x, y)$$

$$\Rightarrow \text{now } u(x, y) = w(\xi(x, y), \eta(x, y))$$

then

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial \xi}{\partial x} + \frac{\partial \eta}{\partial x} \rightarrow \frac{\partial}{\partial x} = \frac{\partial \xi}{\partial x} + \frac{\partial \eta}{\partial x} \\ \frac{\partial u}{\partial y} &= \frac{\partial \xi}{\partial y} + \frac{\partial \eta}{\partial y} \rightarrow \frac{\partial}{\partial y} = \frac{\partial \xi}{\partial y} + \frac{\partial \eta}{\partial y} \\ \frac{\partial^2 u}{\partial x^2} &= \frac{\partial}{\partial x} \left[\frac{\partial \xi}{\partial x} + \frac{\partial \eta}{\partial x} \right] \\ &= \frac{\partial^2 \xi}{\partial x^2} + \frac{\partial \xi}{\partial x} \frac{\partial \eta}{\partial x} + \frac{\partial \eta}{\partial x} \frac{\partial \xi}{\partial x} + \frac{\partial^2 \eta}{\partial x^2} \\ &= \frac{\partial^2 \xi}{\partial x^2} + \frac{\partial \xi}{\partial x} \left[\frac{\partial \eta}{\partial x} + \frac{\partial^2 \eta}{\partial x^2} \right] \\ &\quad + \frac{\partial \eta}{\partial x} \left[\frac{\partial \xi}{\partial x} + \frac{\partial^2 \xi}{\partial x^2} \right] \\ &= \frac{\partial^2 \xi}{\partial x^2} + \left(\frac{\partial \xi}{\partial x} \right)^2 + \frac{\partial \xi}{\partial x} \frac{\partial \eta}{\partial x} + \left(\frac{\partial \eta}{\partial x} \right)^2 \end{aligned}$$

and similarly

$$\begin{aligned} \frac{\partial^2 u}{\partial y^2} &= \frac{\partial}{\partial y} \left(\frac{\partial \xi}{\partial y} + \frac{\partial \eta}{\partial y} \right) + \frac{\partial^2 \xi}{\partial y^2} + \frac{\partial^2 \eta}{\partial y^2} \\ &\quad + \frac{\partial \xi}{\partial y} \frac{\partial \eta}{\partial y} + \frac{\partial \eta}{\partial y} \frac{\partial \xi}{\partial y} \\ \frac{\partial^2 u}{\partial xy} &= \frac{\partial^2 \xi}{\partial xy} + \frac{\partial \xi}{\partial x} \frac{\partial \eta}{\partial y} + \frac{\partial \eta}{\partial x} \frac{\partial \xi}{\partial y} \\ &\quad + 2 \frac{\partial \xi}{\partial x} \frac{\partial \eta}{\partial y} \end{aligned}$$

So the new PDE is

$$\tilde{L}(w) = Aw_{\xi\xi} + 2Bw_{\xi\eta} + Cw_{\eta\eta} + Dw_{\xi} + Ew_{\eta} + Fw = G$$

with

$$A = a\xi_x^2 + 2b\xi_x\xi_y + c\xi_y^2$$

$$B = a\eta_x\xi_x + b(\xi_x\eta_y + \xi_y\eta_x) + c\eta_y\xi_y$$

$$C = a\eta_y^2 + 2b\eta_x\eta_y + c\eta_y^2$$

Another way of writing this is

$$\begin{pmatrix} A & B \\ B & C \end{pmatrix} = \begin{pmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{pmatrix} \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} \xi_x & \eta_x \\ \xi_y & \eta_y \end{pmatrix}$$

$$= J \begin{pmatrix} a & b \\ b & c \end{pmatrix} J^T$$

Now since $\delta(\tilde{x}) = - \begin{vmatrix} a & b \\ b & c \end{vmatrix}$ then

$$\delta(\tilde{x}) = - \begin{vmatrix} A & B \\ B & C \end{vmatrix} = +|J| \delta(x) |J^T| = |J| |J^T| \delta(\tilde{x}) = |J|^2 \delta(\tilde{x})$$

\Rightarrow so provided $|J| \neq 0$ $\delta(\tilde{x})$ has the same sign as $\delta(x)$, as required.

3.3 Canonical forms

We now consider three types of equations:

hyperbolic equations
parabolic equations
and elliptic equations

($\delta(\tilde{x}) > 0$ everywhere)
($\delta(\tilde{x}) = 0$)
($\delta(\tilde{x}) < 0$)

It is possible to find a coordinate transform $(x, y) \rightarrow (\xi, \eta)$ reducing these equations to their canonical forms such that

($\delta(\tilde{x}) = 1/4$) . hyperbolic equations become $u_{\xi\eta} + l_1(u) = g(\xi, \eta)$

($\delta(\tilde{x}) = 0$) . parabolic equations $u_{\xi\xi} + l_1(u) = g(\xi, \eta)$

($\delta(\tilde{x}) = -1$) . elliptic equations $u_{\xi\xi} + u_{\eta\eta} + l_1(u) = g(\xi, \eta)$

where $l_1(u)$ is a linear operator of first order