

The eigenvectors and eigenvalues are

$$\begin{cases} v_n(x) = \sin\left(\frac{n\pi x}{L}\right) \\ \lambda_n = \frac{n^2\pi^2}{L^2} \end{cases} \quad (n=1, \dots, \infty)$$

let's redefine $m = n-1$ so that

$$\begin{cases} \lambda_m = \frac{(m+1)^2\pi^2}{L^2} \\ v_m(x) = \sin\left(\frac{(m+1)\pi x}{L}\right) \end{cases} \quad (m=0, \dots, \infty)$$

let's verify each of the properties again:

(a) Symmetry of the operator

$$\begin{aligned} & \int_0^L \left(\frac{d^2 u}{dx^2} v - \frac{d^2 v}{dx^2} u \right) dx \\ &= \left[v \frac{du}{dx} \right]_0^L - \int_0^L \frac{du}{dx} \frac{dv}{dx} dx \\ & \quad - \left[u \frac{dv}{dx} \right]_0^L + \int_0^L \frac{du}{dx} \frac{dv}{dx} dx \\ &= \left[v \frac{du}{dx} - u \frac{dv}{dx} \right]_0^L = 0 \end{aligned}$$

for u, v satisfying
the same bcs

(b) Orthogonality of the eigenfunctions

assume $n \neq m$:

$$\begin{aligned} \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx &= \int_0^L \left[\cos\left(\frac{(n-m)\pi x}{L}\right) - \cos\left(\frac{(n+m)\pi x}{L}\right) \right] \frac{dx}{2} \\ &= \frac{L}{(n-m)\pi} \left[\sin\left(\frac{(n-m)\pi x}{L}\right) \right]_0^L - \frac{L}{(n+m)\pi} \left[\sin\left(\frac{(n+m)\pi x}{L}\right) \right]_0^L \\ &= 0 \end{aligned}$$

if $n=m$ then

$$\begin{aligned} \int_0^L \sin^2\left(\frac{n\pi x}{L}\right) dx &= \int_0^L \left(1 - \cos\left(\frac{2n\pi x}{L}\right) \right) \frac{dx}{2} \\ &= \frac{L}{2} \end{aligned}$$

(c) λ is indeed real

(e) $\{\lambda_n\}$ form an infinite monotonous sequence

(f) Every function f satisfying $f(0) = f(L) = 0$ can be written as

$$f(x) = \sum_{n=1}^{\infty} a_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\text{with } a_n = \frac{\langle f(x), \sin\left(\frac{n\pi x}{L}\right) \rangle}{\langle \sin\left(\frac{n\pi x}{L}\right), \sin\left(\frac{n\pi x}{L}\right) \rangle}$$

$$= \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx \quad \rightarrow \text{see previous Chapter}$$

⑦

Define $R(u) = - \frac{\int_a^b u \mathcal{L}(u) dx}{\int_a^b r u^2 dx}$

(the Rayleigh quotient)

then, the following theorem holds:

Theorem: • The principal eigenvalue λ_0 of a regular Sturm-Liouville problem is the solution of

$$\lambda_0 = \inf_{u \in V} R(u) \quad (\text{Rayleigh-Ritz formula})$$

where V is the space of all continuous & differentiable functions on (a, b) such that u satisfy the BCs of the Sturm-Liouville problem, and $u \neq 0$ (not the trivial function)

• The function u_0 for which the minimum of $R(u)$ is achieved is the corresponding eigenfunction of the principal eigenvalue

Proof: let $\{\lambda_0 \dots \lambda_n \dots\}$ be the set of all eigenvalues of the S.L. prob. with $\{v_0 \dots v_n \dots\}$ the set of corresponding orthonormal eigenfunctions

then $u = \sum_n a_n v_n(x)$

and $\mathcal{L}(u) = -\sum_n a_n \lambda_n r(x) v_n(x)$.

Then $\int_a^b u \mathcal{L}(u) dx = \int_a^b -\sum_n \sum_m a_n a_m \lambda_n r(x) v_n(x) v_m(x) dx$

modulo some arguments about exchanging \sum and \int

$= -\sum_n \sum_m \int_a^b a_n a_m \lambda_n r(x) v_n(x) v_m(x) dx$

$= -\sum_n a_n^2 \lambda_n$

$\int_a^b r u^2 dx = \int_a^b \sum_n \sum_m a_n a_m v_n(x) v_m(x) r(x) dx$

same $\rightsquigarrow = \sum_n a_n^2$

\rightarrow
PTD

so $R(u) = \frac{\sum_n a_n^2 \lambda_n}{\sum_n a_n^2}$

now given that we know that $\forall n > 0 \lambda_n > \lambda_0$ then

$$R(u) \geq \frac{\lambda_0 \sum_n a_n^2}{\sum_n a_n^2} = \lambda_0$$

To have equality, we would require that $a_n = 0 \forall n > 0$
so that

$$R(u) = \frac{a_0^2 \lambda_0}{a_0^2} = \lambda_0$$

If $u_0 = a_0 v_0$ then u_0 is indeed the eigenfunction corresponding to the principle eigenvalue λ_0 .

Notes ① given that $\int_a^b u x'(u) dx$

$$= \int_a^b u \left[(p(x)u')' + q(x)u \right] dx$$

$$= \int_a^b q(x)u^2 dx + \left[up(x)u' \right]_a^b$$

$$- \int_a^b p(x)u'^2 dx$$

then

$$R(u) = \inf_{u \in V} \left[\frac{\int_a^b (p(x)u'^2 - q(x)u^2) dx - [uu'p]_a^b}{\int_a^b ru^2 dx} \right]$$


This becomes particularly simple for ^{homogeneous} Neumann or Dirichlet conditions: in that case

$$R(u) = \inf_{u \in V} \left[\frac{\int_a^b [p(x)u'^2 - q(x)u^2] dx}{\int_a^b ru^2 dx} \right]$$

$$R(u) = \frac{\int_0^1 u'^2 + x^2 u^2 dx - [u(1)u'(1) - u(0)u'(0)]}{\int_0^1 u^2 dx}$$

so we see that $R(u) \geq 0 \Rightarrow \lambda_0 \geq 0$

To obtain an estimate for λ_0 , consider a trial function $u(x)$ that satisfy the boundary conditions



→ could try $u(x) = A \cos\left(\frac{\pi}{2}x\right)$

or $u(x) = A(1-x^2)$

Using the 2nd option

$$R(u) = \frac{\int_0^1 A^2 (-2x)^2 + x^2 A^2 (1-x^2)^2 dx}{\int_0^1 A^2 (1-x^2)^2 dx}$$

$$= \frac{\int_0^1 (4x^2 + x^2 - 2x^4 + x^6) dx}{\int_0^1 (1 - 2x^2 + x^4) dx}$$

$$= \frac{\frac{5}{3} - \frac{2}{5} + \frac{1}{7}}{1 - \frac{2}{3} + \frac{1}{5}} = \frac{37}{14}$$

$$\Rightarrow 0 \leq \lambda_0 \leq \frac{37}{14}$$

Exercise; try the same procedure with $A \cos\left(\frac{\pi}{2}x\right)$

Note: The solution has $\lambda_0 = 2.597 \dots$ $\frac{37}{14} = 2.64$

② It is actually possible to show that the sequence of eigenvalues of the S.L. problem $(pu_n)' + qu_n = -\lambda_n u_n$ is also the set of all stationary pts of the Rayleigh quotient $R(u)$ over V , and the eigenfunctions are the functions for which this stationary pt is achieved:

$$\lambda_n = R(v_n)$$

This form is often v. useful to determine the sign of λ_0 , and to obtain order-of-magnitude estimates for it

(6.1) Example 1 . Consider the S.L. problem

$$\begin{cases} u'' + \lambda u = 0 \\ u(0) - u'(0) = 0 \\ u(1) + u'(1) = 0 \end{cases} \quad x \in [0, 1]$$

Here, we have a S.L. problem with

$$\begin{cases} p(x) = 1 \\ q(x) = 0 \\ r(x) = 1 \end{cases}$$

$$\text{so } R(u) = \frac{\int_0^1 u'^2 dx - [u(1)u'(1) - u(0)u'(0)]}{\int_0^1 u^2 dx}$$

but $u'(1) = -u(1)$ and $u'(0) = u(0) \Rightarrow [u(1)u'(1) - u(0)u'(0)] = -u(1)^2 - u(0)^2$
 \rightarrow so clearly $R(u) \geq 0$ for all u , which proves that $\lambda_0 \geq 0$.

(6.4) Example 2 Consider the S.L. problem.

$$\begin{cases} u'' + (\lambda - x^2)u = 0 \\ u'(0) = 0 \\ u(1) = 0 \end{cases}$$

We want to find an estimate for λ_0 .

Here, we have the S.L. problem with

$$\begin{cases} p(x) = 1 \\ q(x) = -x^2 \\ r(x) = 1 \end{cases} \quad \text{so}$$

⑧ Consider a regular S.L problem with eigenvalues $\{\lambda_0, \lambda_1, \dots, \lambda_n, \dots\}$ and eigenfunctions $\{v_0, v_1, \dots, v_n, \dots\}$.
 Then v_n has exactly n roots over the interval (a, b)




In particular, v_0 has no node in (a, b)

Remark: This is why the simplest guess for $y(x)$ for estimating λ_0 actually is also the best.

Example:

$$\begin{cases} u'' + \lambda u = 0 \\ u(0) = u(L) = 0 \end{cases}$$

$$\begin{cases} \lambda_n = \frac{\pi^2 (n+1)^2}{L^2} \\ v_n = \sin\left(\frac{\pi x}{L}(n+1)\right) \end{cases} \quad n \geq 0$$



etc...

... : ... obtained in the ...

⑧ Asymptotic ($n \rightarrow \infty$) approximations to the eigenfunctions and eigenvalues of a regular SL problem

For large n , we know that $\lambda_n \rightarrow +\infty$; if this is the case, it is possible to approximate the eigenfunctions by

$$v_n(x) \approx \frac{1}{(r(x)p(x))^{1/4}} \left\{ \alpha \cos \left[\sqrt{\lambda_n} \int_a^x \sqrt{\frac{r(x')}{p(x')}} dx' \right] + \beta \sin \left[\sqrt{\lambda_n} \int_a^x \sqrt{\frac{r(x')}{p(x')}} dx' \right] \right\}$$

(This formula is derived from the WKB approximation (see AMS 212b))

In that case it's easy to see that

$$\sqrt{\lambda_n} \int_a^b \sqrt{\frac{r(x')}{p(x')}} dx' \approx n\pi$$
$$\Rightarrow \lambda_n \approx \left(\frac{n\pi}{\int_a^b \sqrt{\frac{r(x')}{p(x')}} dx'} \right)^2$$

Example

$$\begin{cases} u'' + \lambda u = 0 \\ u(0) = u'(0) \\ u(1) = -u'(1) \end{cases}$$

we saw that $\lambda \geq 0$

This time, let's look for the eigensolutions:

$$u(x) = A \cos \sqrt{\lambda} x + B \sin \sqrt{\lambda} x$$

to satisfy the bcs we calculate

$$u'(x) = -A\sqrt{\lambda} \sin\sqrt{\lambda}x + B\sqrt{\lambda} \cos\sqrt{\lambda}x$$

so

$$\begin{cases} A = B\sqrt{\lambda} \\ A\cos\sqrt{\lambda} + B\sin\sqrt{\lambda} = +A\sqrt{\lambda}\sin\sqrt{\lambda} - B\sqrt{\lambda}\cos\sqrt{\lambda} \end{cases}$$


$$\Rightarrow \begin{cases} \sqrt{\lambda}\cos\sqrt{\lambda} + \sin\sqrt{\lambda} = +\lambda\sin\sqrt{\lambda} - \sqrt{\lambda}\cos\sqrt{\lambda} \\ A = B\sqrt{\lambda} \end{cases}$$

$$\Rightarrow \begin{cases} 2\sqrt{\lambda}\cos\sqrt{\lambda} = (\lambda-1)\sin\sqrt{\lambda} \\ A = B\sqrt{\lambda} \end{cases}$$

$$\Rightarrow \begin{cases} \tan\sqrt{\lambda} = \frac{2\sqrt{\lambda}}{\lambda-1} \\ A = B\sqrt{\lambda} \end{cases}$$

to find λ , we must solve the equation $\tan x = \frac{2x}{x^2-1}$

Graphically with $x > 0$



\Rightarrow looks like

$$x_n = n\pi$$

for large n

$$\Rightarrow \lambda_n \approx n^2\pi^2$$


Check: using the asymptotic formula with

$$r(x) = p(x) = 1 \quad q(x) = 0 \quad \begin{matrix} a=0 \\ b=1 \end{matrix}$$

$$\Rightarrow \lambda_n \approx (n^2\pi^2). \text{ indeed for large } n.$$

Example of application : radial oscillations of a circular drum & Bessel functions

Let's consider a circular drum, assume it is oscillating in an axisymmetric way :



$$u_{tt} = \frac{c^2}{x} \frac{\partial}{\partial x} \left(x \frac{\partial u}{\partial x} \right)$$

$$u(R, t) = 0$$

$$|u(0, t)| < +\infty$$

At time $t=0$, we hit the drum dead center with a stick, giving it a velocity

$$\begin{cases} u(x, 0) = 0 \\ u_t(x, 0) = e^{-x^2/2\sigma^2} \end{cases} \text{ with } \sigma \ll R.$$

Separation of variables : $u(x, t) = A(x) B(t)$

$$\begin{cases} \frac{d^2 B}{dt^2} = -c^2 \lambda B \\ \frac{1}{x} \frac{d}{dx} \left(x \frac{dA}{dx} \right) = -\lambda A \end{cases}$$

Spatial problem : The second equation represents a singular Sturm-Liouville problem with

$$\begin{cases} p(x) = x \\ q(x) = 0 \\ r(x) = x \end{cases}$$

(singular because p & r vanish @ $x=0$)

What are the solutions?

$$\frac{1}{x} \frac{d}{dx} \left(x \frac{dA}{dx} \right) = -\lambda A \quad \Rightarrow \quad x^2 \frac{d^2 A}{dx^2} + x \frac{dA}{dx} + \lambda x^2 A = 0$$